

## Anti-h FDP LL01002 SPTN-5

### Product overview

---

<b>Catalog number</b>	C-40-0024
<b>Specificity</b>	Antibody recognizes human fibrin degradation protein (FDP)
<b>Description</b>	Monoclonal mouse antibody, cultured in vitro under conditions free from animal-derived components.
<b>Product buffer solution</b>	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	18 months from manufacturing at 2–8 °C
<b>Analyte description</b>	Fibrinogen has a molecular weight of 340,000 and is composed of three pairs of peptide chains (A $\alpha$ , B $\beta$ , C $\gamma$ ) connected by disulfide bonds. It transforms into fibrin under the action of thrombin. Both can be degraded into fragments such as X, Y, D, E, and D dimer by plasmin. Fragments X, Y, D, E, etc. are collectively referred to as FDP. FDP is a sensitive indicator of systemic or local intravascular coagulation and fibrinolysis.

### Parameters tested on each lot

---

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	5.0 mg/ml (+/- 10%)
<b>Immunoreactivity</b>	80–120% compared to the reference sample
<b>IEF Profile</b>	5.4 - 6.3
<b>Purity</b>	≥ 95 %

### Kinetic parameters

---

<b>Association rate constant</b>	To Be Determined (TBD)
<b>Dissociation rate constant</b>	TBD
<b>Affinity constant</b>	TBD
<b>Determination method</b>	-
<b>Determination antigen</b>	-



**Cross-reactivities** Not Determined (N/D)

**Epitope** Not Determined (N/D)

**Pair recommendations**

		DETECTION				
		LL01001	LL01002	LL01003	LL01004	LL01005
CAPTURE	LL01001	-	-	-	+	-
	LL01002	+	-	+	+	+
	LL01003	-	+	-	-	-
	LL01004	-	-	-	-	-
	LL01005	-	-	-	-	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

**Antigens tested** -

Product stability	TEMPERATURE, TIME	RESULT
	-70 °C, 21 days	OK
	-20 °C, 21 days	OK
	+4 °C, 21 days	OK
	+35 °C, 21 days	OK
	+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

**Miscellaneous** -

**References** -

